

Aceclofenac (100mg) + Paracetamol (325mg) + Serratiopeptidase (15mg)

## Aceflick SP tablet

**Aceclofenac (100 mg) + Paracetamol (325 mg) +  
Serratiopeptidase (15 mg) Tablet**

---

### Description:

This combination tablet contains Aceclofenac, Paracetamol, and Serratiopeptidase, designed to provide effective relief from pain, inflammation, and swelling while promoting healing in conditions associated with musculoskeletal injuries, arthritis, and other inflammatory disorders. This combination is commonly prescribed for the relief of pain, fever, and inflammation while aiding in tissue repair.

---

### Composition and Mechanism of Action:

1. **Aceclofenac (100 mg):** A nonsteroidal anti-inflammatory drug (NSAID) that works by inhibiting cyclooxygenase (COX) enzymes, leading to a reduction in the production of prostaglandins, which are chemicals responsible for causing inflammation, pain, and fever. Aceclofenac helps in reducing pain and swelling in conditions like arthritis, sprains, and other inflammatory disorders.
2. **Paracetamol (325 mg):** An analgesic (pain reliever) and antipyretic (fever reducer) that works by inhibiting prostaglandin production in the brain, helping to alleviate pain and reduce fever. It is often used in combination with other medications to enhance pain relief.
3. **Serratiopeptidase (15 mg):** An enzyme that has anti-inflammatory and proteolytic properties, aiding in the breakdown of non-living tissue, reducing swelling, and promoting the healing of injured tissues. Serratiopeptidase helps in reducing the inflammation and swelling associated with conditions like arthritis, injuries, and post-surgical recovery.

### Indications:

This combination tablet is indicated for:

- Pain and inflammation associated with musculoskeletal conditions like arthritis, sprains, strains, and back pain
  - Post-surgical recovery to reduce swelling and promote tissue healing
  - Acute pain management, including soft tissue injuries, joint pain, and inflammatory conditions
  - Symptomatic relief from fever and discomfort associated with infections or injuries
- 

### Dosage and Administration:

- **Adults:** One tablet every 8–12 hours as required, or as directed by your healthcare provider. Do not exceed three tablets in a 24-hour period.
- **Route of Administration:** Oral, to be taken with water.
- **Timing:** Preferably after meals to reduce the risk of stomach irritation.

**Note:** Dosage may vary depending on the severity of the symptoms and the patient's response to treatment. Always follow the instructions of your healthcare provider.

---

### Precautions and Warnings:

1. **Alcohol:** Avoid alcohol while taking this medication as it may increase the risk of liver damage (due to Paracetamol) and stomach irritation (due to Aceclofenac).
2. **Pregnancy and Breastfeeding:** Use during pregnancy or breastfeeding should be done only if prescribed by a doctor, particularly in the third trimester.

3. **Pre-existing Conditions:** Use with caution in individuals with liver or kidney problems, gastrointestinal disorders (e.g., peptic ulcers), or heart disease.
4. **Allergic Reactions:** Discontinue use if signs of an allergic reaction occur (e.g., skin rash, swelling, difficulty breathing).
5. **Renal and Hepatic Health:** Prolonged use of NSAIDs like Aceclofenac may affect kidney function, so regular monitoring is recommended.

- **Other NSAIDs:** Avoid combining with other NSAIDs or corticosteroids due to an increased risk of gastrointestinal irritation or ulcers.
- **Diuretics and Antihypertensives:** Concomitant use with diuretics or blood pressure medications may reduce their effectiveness.
- **Alcohol and Hepatotoxic Drugs:** Combining Paracetamol with alcohol or liver-toxic medications may increase the risk of liver damage.

**Note:** Always inform your healthcare provider about any other medications, including over-the-counter drugs and supplements, to avoid potential interactions.

---

## Contraindications:

- Known hypersensitivity or allergy to Aceclofenac, Paracetamol, Serratiopeptidase, or other NSAIDs
- Active gastrointestinal bleeding, peptic ulcer disease, or severe liver impairment
- Severe renal impairment or heart failure
- Pregnancy (especially in the third trimester) or breastfeeding without medical supervision

---

## Storage:

- Store in a cool, dry place at room temperature, away from direct sunlight and moisture.
- Keep out of reach of children.

---

## Side Effects:

- **Common Side Effects:** Nausea, vomiting, abdominal discomfort, indigestion, dizziness, headache, and mild gastrointestinal disturbances.
- **Serious Side Effects (rare):** Gastrointestinal bleeding (e.g., black stools, vomiting blood), signs of liver damage (yellowing of skin or eyes), allergic reactions (rash, swelling, difficulty breathing), and kidney dysfunction (e.g., decreased urine output).

If any serious side effects occur, stop using the medication and seek immediate medical attention.

---

## Disclaimer:

This medication should be used only under medical supervision. It is intended for symptomatic relief and does not address the underlying cause of pain or inflammation. Always follow your healthcare provider's instructions for proper use, dosage, and duration of treatment.

Manufactured in India for:

**Cafoli**<sup>TM</sup>  
L I F E C A R E

**Cafoli Lifecare Pvt. Ltd.**

**(An ISO 9001: 2015 Certified Co.)**

Plot no.: 367-FF, Industrial Area Phase-I,  
Panchkula-134113

TM: Trademark Applied for

---

## Drug Interactions:

- **Anticoagulants:** Aceclofenac may increase the risk of bleeding when taken with blood thinners like warfarin.