

Cabstyle 0.5 Tablet

Cabstyle 0.5 Tablet – Cabergoline 0.5mg for Hyperprolactinemia Treatment

Composition:

Each **Cabstyle 0.5 Tablet** contains:

- **Cabergoline 0.5 mg** – A dopamine agonist that reduces prolactin levels by stimulating dopamine receptors in the pituitary gland.

Description:

Cabstyle 0.5 Tablet is an **ergoline-derived dopamine agonist** used primarily for the **treatment of hyperprolactinemia**, a condition characterized by excessive prolactin hormone levels. By acting on **dopamine D2 receptors**, Cabergoline effectively inhibits prolactin secretion, restoring normal hormonal balance.

Pharmacological Action:

- **Reduces excessive prolactin production**, treating menstrual irregularities, infertility, and lactation disorders.
- Used in **Parkinson's disease** as an adjunct therapy due to its dopamine-stimulating effects.
- Helps prevent **lactation (milk production) in postpartum women** when suppression is necessary.

Indications:

Cabstyle 0.5 Tablet is recommended for:

- **Hyperprolactinemia-related disorders** such as irregular menstrual cycles, infertility, and galactorrhea.
- **Suppression of lactation** in postpartum women when breastfeeding is not desired.
- **Adjunct therapy in Parkinson's disease** (in some cases).

Dosage & Administration:

- **Hyperprolactinemia: 0.25mg to 0.5mg twice weekly**, adjusted as needed.
- **Lactation Suppression: 1mg as a single dose on the first postpartum day.**
- **Take with food to reduce gastrointestinal side effects.**

Contraindications:

- **Uncontrolled hypertension or cardiovascular disorders.**
- **History of pulmonary, pericardial, or retroperitoneal fibrotic disorders.**
- **Severe hepatic impairment.**
- **Pregnant women unless prescribed by a doctor.**

Warnings & Precautions:

- **Regular monitoring of heart function** is advised, as Cabergoline may cause **cardiac valvulopathy**.
- **Use caution in patients with psychiatric disorders**, as it may cause mood changes.
- Avoid **alcohol and sedatives** while taking this medication.

Adverse Effects:

- **Nausea, vomiting, dizziness, and fatigue.**
- **Postural hypotension (low blood pressure upon standing).**
- **Sleep disturbances or hallucinations** (rare but possible).

Drug Interactions:

- Avoid concurrent use with **dopamine antagonists** (e.g., antipsychotics) as they may reduce effectiveness.
- **Macrolide antibiotics (e.g., erythromycin)** can increase Cabergoline levels in the body.
- **Antihypertensive drugs** may enhance the risk of low blood pressure.

Storage Conditions:

- Store at **room temperature (below 30°C)**, away from moisture.
- Keep out of reach of **children**.

Conclusion:

Cabstyle 0.5 Tablet (Cabergoline 0.5mg) is a **clinically proven dopamine agonist** for **hyperprolactinemia and lactation suppression**, ensuring effective hormonal regulation and reproductive health benefits.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,
Panchkula-134113

TM: Trademark Applied for