

# Chymoedge Forte Tablet

## Product Information: Trypsin-Chymotrypsin (100,000 AU) Tablet

---

### Composition:

#### Each tablet contains:

- **Trypsin-Chymotrypsin: 100,000 AU (Activity Units).**
- 

### Pharmacological Class:

- **Enzymes: Proteolytic enzyme combination.**
- 

### Indications:

#### The Trypsin-Chymotrypsin 100,000 AU Tablet is indicated for:

1. **Inflammation and Edema:**
  - Management of post-operative or post-traumatic swelling and inflammation.
2. **Soft Tissue Injuries:**
  - Reduces swelling and pain in conditions like sprains, strains, and muscle injuries.
3. **Respiratory Conditions:**
  - Adjunct therapy for clearing excessive mucus in respiratory infections.
4. **Wound Healing:**
  - Supports tissue repair and accelerates healing in injuries or ulcers.

#### 5. Arthritis and Joint Pain:

- Helps alleviate inflammation in osteoarthritis or rheumatoid arthritis.
- 

### Mechanism of Action:

- **Proteolytic Activity:** Trypsin and chymotrypsin are enzymes that break down proteins into smaller peptides. They reduce inflammation by hydrolyzing inflammatory mediators, which helps decrease edema, improve blood flow, and support tissue repair.
  - **Anti-Inflammatory Effect:** These enzymes modulate inflammatory processes and improve healing by clearing necrotic tissue, enhancing the availability of essential nutrients and oxygen.
- 

### Dosage and Administration:

- **Recommended Dosage:** 1–2 tablets, 2–3 times daily.
  - **Administration:**
    - Take orally with water, preferably on an empty stomach or as advised by your healthcare provider.
    - Swallow the tablet whole; do not crush or chew.
- 

### Contraindications:

- **Hypersensitivity:** Allergy to trypsin, chymotrypsin, or any other component in the formulation.
  - **Active Bleeding Disorders:** Conditions like hemophilia or active peptic ulcers.
  - **Severe Liver or Kidney Disease:** Use with caution in patients with significant hepatic or renal impairment.
- 

### **Warnings and Precautions:**

1. **Pregnancy and Lactation:**
    - Use only under medical supervision during pregnancy or breastfeeding.
  2. **Bleeding Risk:**
    - Avoid in individuals with coagulation disorders or those taking anticoagulants (e.g., warfarin, aspirin) due to the potential for increased bleeding.
  3. **Surgery:**
    - Discontinue before surgery as advised by a physician.
  4. **Gastrointestinal Disorders:**
    - Use cautiously in individuals with a history of gastric or duodenal ulcers.
- 

### **Adverse Effects:**

- **Common Side Effects:**
    - Nausea or gastrointestinal discomfort.
    - Mild allergic reactions (e.g., rash or itching).
  - **Rare but Serious Side Effects:**
    - Severe allergic reactions, such as swelling or difficulty breathing.
    - Prolonged bleeding or hematoma formation.
- 

### **Drug Interactions:**

- **Anticoagulants (e.g., warfarin, heparin):**
    - Increased risk of bleeding; use with caution.
  - **Antiplatelet Agents (e.g., aspirin, clopidogrel):**
    - May enhance the anticoagulant effect, raising the bleeding risk.
  - **Anti-inflammatory Medications:**
    - May be used concomitantly to enhance anti-inflammatory benefits but only under medical guidance.
- 

### **Use in Special Populations:**

- **Pregnancy:** Use with caution; consult a healthcare provider.
  - **Lactation:** Safety during breastfeeding has not been fully established.
  - **Pediatrics:** Safety and efficacy in children should be determined by a physician.
  - **Elderly:** Well-tolerated; dosage adjustments may not be necessary unless there are underlying health conditions.
- 

### **Storage:**

- Store in a cool, dry place, below 25°C.
  - Protect from moisture and direct sunlight.
  - Keep out of reach of children.
- 

### **Packaging:**

- Available in strips of 10 tablets.
  - Typically packed in boxes containing 10 strips.
-

---

## **Patient Instructions:**

- 1. Follow the prescribed dose by your healthcare provider; do not self-medicate.**
  - 2. Avoid alcohol during treatment to minimize gastrointestinal side effects.**
  - 3. Report any unusual symptoms (e.g., prolonged bleeding, allergic reactions) to your doctor immediately.**
  - 4. Do not skip doses; take at regular intervals for optimal effectiveness.**
- 

**Trypsin-Chymotrypsin (100,000 AU) Tablet is a potent anti-inflammatory and proteolytic enzyme supplement widely used to reduce inflammation, promote healing, and alleviate pain associated with injuries, surgeries, and other inflammatory conditions. Always use under medical supervision for safe and effective results.**

Manufactured in India for:



**Cafoli Lifecare Pvt. Ltd.**

**(An ISO 9001: 2015 Certified Co.)**

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for