

Ciproplit TZ Tablet

Ciprofloxacin (500 mg) + Tinidazole (600 mg) Tablet

Composition

Each tablet contains:

- Ciprofloxacin – 500 mg
 - Tinidazole – 600 mg
-

Description

This combination tablet contains Ciprofloxacin, a fluoroquinolone antibiotic, and Tinidazole, a nitroimidazole antimicrobial. The combination is used to treat infections caused by both aerobic and anaerobic bacteria, providing a broad-spectrum antimicrobial effect.

- **Ciprofloxacin:** A broad-spectrum antibiotic that works by inhibiting bacterial DNA synthesis through the inhibition of DNA gyrase. It is effective against a wide range of Gram-negative and Gram-positive bacteria.
 - **Tinidazole:** A nitroimidazole antibiotic that works by interfering with the DNA synthesis of anaerobic microorganisms, effectively treating infections caused by anaerobic bacteria and some protozoa.
-

Indications

This combination is indicated for the treatment of infections caused by susceptible organisms, including:

- Gastrointestinal infections (e.g., bacterial gastroenteritis, giardiasis, amebiasis)
 - Intra-abdominal infections (e.g., peritonitis, abscesses)
 - Pelvic inflammatory disease (PID)
 - Urinary tract infections (UTIs)
 - Bacterial vaginosis
 - Gynecological infections
-

Dosage and Administration

- **Adults:**
 - **For Infections (General):** The typical dosage is 1 tablet (500 mg Ciprofloxacin + 600 mg Tinidazole) once or twice daily for 7 to 10 days, depending on the severity and location of the infection.
 - **For Pelvic Inflammatory Disease:** The recommended dosage is 1 tablet (500 mg Ciprofloxacin + 600 mg Tinidazole) twice daily for 14 days.
 - **For Urinary tract infections:** The typical dosage is 1 tablet (500 mg Ciprofloxacin + 600 mg Tinidazole) once or twice a day for 7–10 days.
- **Children:** This combination should not be used in children under the age of 12 unless specifically prescribed by a doctor, as the safety and efficacy

in pediatric patients have not been established for this combination.

- **Elderly:** Dosage may need to be adjusted for elderly patients, especially those with renal or hepatic impairment.

Administration:

- Take the tablet whole with a full glass of water.
- The tablet can be taken with or without food, but it should be taken at the same time each day for maximum effectiveness.
- Do not chew or crush the tablet.

Warnings and Precautions

- **Tendon Damage:** Ciprofloxacin has been associated with an increased risk of tendonitis and tendon rupture, especially in older adults or those using corticosteroids. Discontinue if you experience tendon pain or swelling.
- **Central Nervous System (CNS) Effects:** Ciprofloxacin can cause dizziness, seizures, or confusion, particularly in elderly patients. If any CNS symptoms occur, consult your doctor immediately.

Side Effects

Common side effects may include:

- Nausea, vomiting, or diarrhea
- Headache
- Dizziness or lightheadedness
- Abdominal pain or discomfort
- Rash
- Metallic taste (due to Tinidazole)

Serious side effects (rare but potentially severe):

- Severe allergic reactions (anaphylaxis, swelling, difficulty breathing)
- Blood disorders (low white blood cell count, low platelet count)
- Severe gastrointestinal disturbances (e.g., persistent diarrhea, colitis)

Seek immediate medical attention if you experience any of these serious side effects.

- of action, as resistance may develop.

Storage

- Store at room temperature, below 25°C, in a cool, dry place, away from direct sunlight and moisture.
- Keep out of reach of children.

Note: Always follow your healthcare provider's instructions for the correct dosage and duration of therapy. Do not take two doses at once if you miss a dose. If you experience any side effects or the condition does not improve, contact your doctor.

Manufactured in India for:

CafoliTM
L I F E C A R E

Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for