

Amoxicillin (750mg) + Pantoprazole (40mg) + Clarithromycin (500mg)

Panpital HP Kit

Amoxicillin (750 mg) + Pantoprazole (40 mg) + Clarithromycin (500 mg) Tablet

Description:

Amoxicillin (750 mg), Pantoprazole (40 mg), and Clarithromycin (500 mg) combination tablet is a multi-drug therapy commonly used in the treatment of *Helicobacter pylori* infection and other related gastrointestinal disorders. This combination helps eradicate H.

Composition:

- Amoxicillin: 750 mg
 - Pantoprazole: 40 mg
 - Clarithromycin: 500 mg
-

Indications:

This tablet is indicated for:

1. **Helicobacter pylori Infection:** Used as part of a triple therapy regimen to treat H. pylori infection associated with peptic ulcers or gastritis.
2. **Peptic Ulcers:** To aid in the healing of ulcers by eradicating the causative bacterial infection and reducing stomach acid.
3. **Gastroesophageal Reflux Disease (GERD):** Helps in managing symptoms of acid reflux while treating any bacterial component present in the stomach.

4. **Gastritis:** For treatment of gastric inflammation caused by bacterial infections, particularly H. pylori.
-

Dosage and Administration:

- **Usual Adult Dose:**
 - Take one tablet once or twice daily, as prescribed by a healthcare provider, usually for 7 to 14 days, depending on the severity of the infection and treatment guidelines.
 - The tablet should be swallowed whole, with or without food, and should not be crushed or chewed.
-

Side Effects:

- **Common:**
 - Diarrhea, nausea, abdominal pain, flatulence, or headache.
 - Taste disturbances or a dry mouth may occur.
- **Serious but Rare:**
 - Allergic reactions such as rash, itching, or swelling, particularly in sensitive individuals.
 - Gastrointestinal bleeding: Black, tarry stools or vomiting blood (signs of gastrointestinal bleeding).
 - Severe liver reactions: Jaundice, dark urine, or abdominal pain (due to

Pantoprazole or Clarithromycin).

- **Heart-related issues:** Chest pain, irregular heartbeat (especially with high doses or pre-existing heart conditions).
- **Antibiotic resistance:** Use only as prescribed to avoid the development of antibiotic-resistant bacteria.

your doctor of all medications you are currently taking.

Storage:

- **Store at room temperature (15-30°C), away from direct light and moisture.**
- **Keep the tablet in its original packaging and out of the reach of children.**

Precautions:

- 1. Allergic Reactions:** If you have a history of allergies to penicillin, macrolides, or PPIs, inform your healthcare provider before using this combination.
- 2. Liver Function:** Use with caution in individuals with liver disease, especially when taking Clarithromycin or Pantoprazole. Liver function should be monitored during treatment.
- 3. Renal Impairment:** Caution is required in patients with impaired kidney function.
- 4. Pregnancy and Breastfeeding:** Consult a healthcare provider before using during pregnancy or breastfeeding, as some of the components may pass into breast milk.
- 5. Drug Interactions:**
 - **Clarithromycin may interact with other medications, including statins, warfarin, and certain antiepileptic drugs.**
 - **Pantoprazole can interact with drugs such as clopidogrel, methotrexate, and certain antifungal medications. Inform**

Note:

This medication should be used exactly as prescribed to ensure effective treatment and avoid complications. Do not skip doses, and ensure you complete the entire treatment course. If you experience any serious side effects, such as allergic reactions, severe abdominal pain, or signs of an infection, contact your doctor immediately.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for