

## Clipsil O Suspension

### Magaldrate I.P. 400mg + Simethicone I.P. 20mg Suspension (Mango & Mint Flavour)

#### Description:

This suspension is a combination of **Magaldrate**, an antacid that neutralizes excess stomach acid, and **Simethicone**, an anti-flatulent that helps relieve gas and bloating. The dual-action formula provides fast and effective relief from acidity, indigestion, heartburn, and gas-related discomfort. The pleasant **Mango & Mint** flavour enhances palatability, making it easier to consume.

#### Pharmacological Action:

##### 1. Magaldrate

- Acts as an acid-neutralizing agent by converting into magnesium and aluminum hydroxide in the stomach.
- Provides a protective coating to the stomach lining, preventing irritation from gastric acid.
- Helps in the management of conditions like gastritis, peptic ulcers, and acid reflux.

##### 2. Simethicone

- Works by breaking down gas bubbles in the stomach and intestines, making them easier to eliminate.
- Provides relief from bloating, discomfort, and abdominal distension.
- Does not affect stomach acid production, making it a safe choice for gas-related symptoms.

#### Indications:

This suspension is indicated for the symptomatic relief of:

- Hyperacidity and acid reflux
- Heartburn and indigestion

- Gastric irritation and bloating
- Flatulence and gas-related discomfort
- Peptic ulcers and gastritis

#### Dosage & Administration:

- **Adults:** 5-10ml after meals or as directed by a physician.
- **Children:** Use under medical supervision.
- Shake well before use.
- Do not exceed the recommended dose.

#### Contraindications:

- Hypersensitivity to Magaldrate, Simethicone, or any other ingredients in the formulation.
- Patients with severe kidney disease or electrolyte imbalances should consult a doctor before use.

#### Warnings & Precautions:

- Long-term use may lead to an imbalance in electrolytes and should be taken under medical supervision.
- Patients with a history of kidney disease should use with caution due to the presence of magnesium and aluminum compounds.
- Avoid consuming alcohol or caffeine-rich beverages as they can increase acid production.

#### Side Effects:

Although generally well-tolerated, some individuals may experience:

- Mild diarrhea or constipation
- Nausea or stomach cramps
- Allergic reactions in rare cases

If any severe side effects occur, discontinue use and seek medical attention.

---

### Drug Interactions:

- May interfere with the absorption of certain medications like antibiotics (tetracyclines, fluoroquinolones), iron supplements, and thyroid medications.
  - It is advisable to take other medications at least **1 to 2 hours before or after** this suspension.
- 

### Storage Instructions:

- Store in a **cool and dry place** below **25°C**.
  - Keep the bottle tightly closed when not in use.
  - Protect from **direct sunlight and moisture**.
  - Keep **out of reach of children**.
- 

### Presentation:

Available in a **[specify volume]** bottle with a measuring cup for accurate dosage.

For **therapeutic use only**. Should be used under **medical supervision**.

Manufactured in India for:



**Cafoli Lifecare Pvt. Ltd.**

**(An ISO 9001: 2015 Certified Co.)**

Plot no.: 367-FF, Industrial Area Phase-I,  
Panchkula-134113

TM: Trademark Applied for