# **Coldstave P Suspension**

Chlorpheniramine Maleate (1 mg) + Paracetamol (125 mg) + Phenylephrine (5 mg) Suspension

## **Description**

This combination suspension contains Chlorpheniramine Maleate, Paracetamol, and Phenylephrine, which work synergistically to relieve symptoms of common cold, flu, and allergies. It provides relief from fever, nasal congestion, and allergic reactions while soothing discomfort associated with upper respiratory tract infections.

#### **Composition**

## Each 5 mL contains:

- Chlorpheniramine Maleate: 1 mg (Antihistamine)
- Paracetamol: 125 mg (Analgesic and Antipyretic)
- Phenylephrine Hydrochloride: 5 mg (Decongestant)

## **Excipients: Q.S.**

## **Pharmacological Action**

- 1. Chlorpheniramine Maleate: A histamine H1 receptor antagonist that reduces allergic symptoms like sneezing, runny nose, and itching.
- 2. Paracetamol: An analgesic and antipyretic that helps relieve fever, headaches, and minor aches and pains.
- 3. Phenylephrine Hydrochloride: A sympathomimetic decongestant that reduces nasal swelling and congestion by constricting blood vessels in the nasal passage.

#### **Indications**

#### This suspension is indicated for:

- Common cold and flu symptoms.
- Nasal congestion.
- Allergic rhinitis.
- Headache and mild fever.
- Sneezing, runny nose, and watery eyes.

#### **Dosage and Administration**

- Children (6-12 years): 5 mL every 6-8 hours or as directed by a physician.
- Children (2-6 years): 2.5 mL every 6-8 hours or as directed by a physician.
- Below 2 years: Consult a pediatrician before use.

Shake well before use. Use a calibrated measuring device to ensure accurate dosing.

#### **Contraindications**

- Hypersensitivity to any of the ingredients.
- Severe hypertension or coronary artery disease.
- Patients on monoamine oxidase inhibitors (MAOIs).

#### **Precautions**

- Use with caution in patients with liver or kidney impairment.
- Avoid concomitant use with other paracetamol-containing products to prevent overdose.
- Do not exceed the recommended dose.

#### **Adverse Effects**

- Drowsiness (due to Chlorpheniramine Maleate).
- Nausea, vomiting, or upset stomach.
- Mild dizziness or headache.
- Allergic reactions (rare).

If any severe adverse effects occur, discontinue use and consult a doctor immediately.

## **Storage Instructions**

- Store at a temperature not exceeding 30°C.
- Protect from light and moisture.
- Keep out of reach of children.

## **Presentation**

Available in bottles of 60 mL / 100 mL / 120 mL.

Note: This is a symptomatic relief medicine. Consult a doctor if symptoms persist for more than 5 days.

Manufactured in India for:



(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for