# Ambroxol (7.5mg) + Guaifenesin (12.5mg) + Terbutaline (0.25mg)

# **Cufsniff AM Drop**

Ambroxol (7.5 mg) + Guaifenesin (12.5 mg) + Terbutaline (0.25 mg) Drops

# **Description**

This combination drop is formulated to provide effective relief from respiratory conditions associated with productive cough, chest congestion, and bronchospasms. It contains Ambroxol, Guaifenesin, and Terbutaline, which work synergistically to improve mucus clearance, ease breathing, and relieve bronchial inflammation.

## Composition

## Each mL contains:

- Ambroxol Hydrochloride: 7.5 mg (Mucolytic)
- Guaifenesin: 12.5 mg (Expectorant)
- Terbutaline Sulfate: 0.25 mg (Bronchodilator)

**Excipients: Q.S.** 

## **Pharmacological Action**

- 1. Ambroxol Hydrochloride: Acts as a mucolytic agent, reducing the viscosity of mucus and aiding its expulsion from the airways.
- 2. Guaifenesin: An expectorant that helps thin mucus, making it easier to cough out and relieving chest congestion.
- 3. Terbutaline Sulfate: A bronchodilator that relaxes the smooth muscles of the airways, alleviating bronchospasms and improving airflow.

#### **Indications**

## This drop is indicated for:

- Respiratory conditions with productive cough.
- Bronchitis and chest congestion.
- Bronchospasm associated with asthma or other obstructive airway diseases.

## **Dosage and Administration**

- Infants (up to 1 year): 5-10 drops 2-3 times daily, or as directed by a physician.
- Children (1-5 years): 10-15 drops 2-3 times daily, or as directed by a physician.

Shake well before use. Use a calibrated dropper for accurate dosing. Administer orally after meals.

#### **Contraindications**

- Hypersensitivity to any of the ingredients.
- Patients with severe hypertension or cardiac arrhythmias.
- Use in premature infants without medical supervision.

#### **Precautions**

- Use cautiously in patients with cardiovascular disorders, hyperthyroidism, or diabetes.
- Do not exceed the prescribed dose.
- If symptoms persist or worsen, consult a pediatrician.
- Not intended for long-term use unless prescribed.

## **Adverse Effects**

- Common: Mild nausea, vomiting, or abdominal discomfort.
- Less common: Dry mouth, dizziness, or palpitations.
- Rare: Allergic reactions such as rash, swelling, or difficulty breathing.

Discontinue use and seek medical attention if adverse reactions occur.

## **Storage Instructions**

- Store below 30°C in a dry place.
- Protect from light and moisture.
- · Keep out of reach of children.

#### **Presentation**

Available in 15 mL / 30 mL bottles with a calibrated dropper.

Note: This medication is for symptomatic relief and should be used under medical supervision. If the condition persists or worsens, seek medical advice.

Manufactured in India for:



(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for