

Ambroxol (15mg/5ml) + Guaifenesin (50mg/5ml) + Menthol (2.5mg/5ml) + Terbutaline (1.25mg/5ml)

Cufsniff AM Syrup

Ambroxol (15 mg/5 mL) + Guaifenesin (50 mg/5 mL) + Menthol (2.5 mg/5 mL) + Terbutaline (1.25 mg/5 mL) Syrup

Description

This combination syrup provides effective relief from cough, chest congestion, and bronchospasm, typically seen in conditions like asthma, bronchitis, and other respiratory tract infections. The combination of Ambroxol, Guaifenesin, Menthol, and Terbutaline works synergistically to facilitate mucus clearance, reduce bronchospasms, soothe throat irritation, and improve breathing.

Composition

Each 5 mL of syrup contains:

- **Ambroxol Hydrochloride: 15 mg (Mucolytic)**
- **Guaifenesin: 50 mg (Expectorant)**
- **Menthol: 2.5 mg (Soothing agent)**
- **Terbutaline Sulfate: 1.25 mg (Bronchodilator)**

Excipients: Q.S.

Pharmacological Action

1. **Ambroxol Hydrochloride: Breaks down thick, sticky mucus, making it easier to expel from the airways, improving lung function.**
2. **Guaifenesin: Thins the mucus, promoting its clearance and reducing chest congestion.**
3. **Menthol: Provides a cooling effect and soothes throat irritation, helping to**

relieve coughing and irritation in the respiratory tract.

4. **Terbutaline Sulfate: Relaxes bronchial smooth muscles, dilates the airways, and improves airflow by reducing bronchospasms.**

Indications

This syrup is indicated for the symptomatic relief of:

- **Productive cough associated with respiratory tract infections, including bronchitis.**
- **Asthma and other obstructive pulmonary diseases with bronchospasms.**
- **Chest congestion and difficulty clearing mucus.**
- **Throat irritation caused by excessive coughing.**

Dosage and Administration

- **Children (2-6 years): 2.5 mL (½ teaspoon) 2-3 times daily, or as directed by a physician.**
- **Children (6-12 years): 5 mL (1 teaspoon) 2-3 times daily, or as directed by a physician.**
- **Adults and Adolescents (12 years and above): 10 mL (2 teaspoons) 2-3 times daily, or as directed by a physician.**

Shake well before use. Administer orally after meals.

Contraindications

- Hypersensitivity to any of the components.
 - Severe hypertension, ischemic heart disease, or tachyarrhythmias.
 - Children under 2 years of age, unless prescribed by a healthcare professional.
-

Precautions

- Use cautiously in patients with cardiovascular disorders, diabetes, or hyperthyroidism.
 - Avoid exceeding the prescribed dosage.
 - Consult a doctor if symptoms persist for more than 5 days or worsen.
 - Not recommended for prolonged use unless advised by a healthcare professional.
 - If you experience dizziness, palpitations, or difficulty breathing, discontinue use and consult a doctor immediately.
-

Adverse Effects

- **Common:** Nausea, mild abdominal discomfort, dizziness, dry mouth, or throat irritation.
- **Less common:** Headache, tremors, or palpitations.
- **Rare:** Allergic reactions such as rash, itching, swelling, or difficulty breathing.

In case of severe adverse reactions, seek immediate medical attention.

Storage Instructions

- Store below 30°C in a dry place.
 - Protect from light and moisture.
 - Keep out of reach of children.
-

Presentation

Available in 100 mL / 200 mL bottles with a measuring cap for accurate dosing.

Note: This syrup provides symptomatic relief and should be used under the guidance of a healthcare provider. For best results, adhere strictly to the prescribed dosage and frequency.

Manufactured in India for:

CafoliTM
L I F E C A R E

Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,
Panchkula-134113

TM: Trademark Applied for