

DecplisT K Tablet

Diclofenac (50 mg) + Serratiopeptidase (10 mg) Tablet

Description

This combination tablet contains Diclofenac, a nonsteroidal anti-inflammatory drug (NSAID), and Serratiopeptidase, a proteolytic enzyme, to provide enhanced relief from pain and inflammation. It is used in the treatment of musculoskeletal pain, inflammation, and other conditions associated with pain and swelling, such as arthritis, sprains, and post-surgical pain.

Composition

Each tablet contains:

- **Diclofenac Sodium: 50 mg** (Nonsteroidal anti-inflammatory drug)
- **Serratiopeptidase: 10 mg** (Proteolytic enzyme)

Excipients: Q.S.

Pharmacological Action

1. **Diclofenac Sodium:** Inhibits cyclooxygenase (COX), leading to a decrease in the production of prostaglandins, which are responsible for causing pain, fever, and inflammation. It effectively reduces pain, swelling, and stiffness associated with various inflammatory conditions.
2. **Serratiopeptidase:** An enzyme that helps to break down protein molecules, promoting the resolution of inflammation and reducing swelling. It aids in the faster absorption and action of diclofenac while helping to alleviate pain and improve mobility.

Indications

This combination tablet is indicated for the treatment of:

- **Acute pain:** Due to injury, sprains, strains, and trauma.
- **Chronic pain:** Associated with osteoarthritis, rheumatoid arthritis, and ankylosing spondylitis.
- **Post-surgical pain:** Following orthopedic, dental, or other surgical procedures.
- **Inflammatory conditions:** Such as tendonitis, bursitis, and soft tissue inflammation.

Dosage and Administration

- **Adults:**
 - 1 tablet taken once or twice daily, preferably with food to reduce the risk of gastrointestinal irritation.
 - The dose may be adjusted by the healthcare provider based on the severity of pain and inflammation.
 - Maximum daily dose: Do not exceed 100 mg of Diclofenac per day.
- **Administration:** Swallow the tablet whole with a glass of water. Do not crush or chew the tablet.

Contraindications

- Hypersensitivity to Diclofenac, Serratiopeptidase, or any other components of the tablet.
- Active gastrointestinal bleeding or peptic ulcers.
- Severe liver or kidney impairment.

- History of severe heart failure, or active cardiovascular disease.
- Pregnant women, particularly during the third trimester, and breastfeeding mothers (unless prescribed by a healthcare professional).

- Store at room temperature (15-30°C), away from direct sunlight and moisture.
- Keep the tablet in its original packaging to protect it from light and moisture.
- Keep out of the reach of children.

Precautions

- **Caution in Cardiovascular Conditions:** Use with caution in patients with a history of heart disease, hypertension, or stroke.
- **Gastrointestinal Care:** May cause gastrointestinal irritation, ulceration, or bleeding; administer with food to minimize this risk.
- **Renal and Hepatic Dysfunction:** Periodically monitor liver and kidney function in long-term use.
- **Elderly Patients:** Increased risk of side effects, especially gastrointestinal issues and cardiovascular effects.
- **Pregnancy:** Not recommended in the third trimester of pregnancy. Use in the first and second trimester should only be considered if the benefits outweigh the risks.

Presentation

- Available in blister packs containing 10 tablets or 20 tablets.

Note: This medication should be used as directed by a healthcare provider. Do not exceed the prescribed dose, and consult a doctor if symptoms persist or worsen.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,
Panchkula-134113

TM: Trademark Applied for

Adverse Effects

- **Common:** Nausea, abdominal pain, indigestion, headache, dizziness, or mild gastrointestinal discomfort.
- **Less common:** Diarrhea, constipation, skin rash, or fluid retention.
- **Serious:** Gastrointestinal bleeding, ulcers, kidney or liver damage, cardiovascular events (e.g., heart attack, stroke).

Seek immediate medical attention if you experience: severe abdominal pain, black or bloody stools, chest pain, shortness of breath, or swelling of the extremities.

Storage Instructions