

Information For the User
Diclofenac (75mg/ml) Injection

Decplis[®] Injection

Brand Name:
Decplis Injection

Composition:

Each ml contains:

Diclofenac Sodium I.P. ... 75 mg

Water for Injection I.P. ... q.s.

Dosage Form:

Injection for intramuscular or slow intravenous administration.

Pharmacology:

Diclofenac sodium is a non-steroidal anti-inflammatory drug (NSAID) that exhibits potent analgesic, anti-inflammatory, and antipyretic properties. It works primarily by inhibiting cyclooxygenase (COX-1 and COX-2) enzymes, thereby reducing the synthesis of prostaglandins responsible for pain, inflammation, and fever.

Indications:

- Relief of acute musculoskeletal pain
- Postoperative pain management
- Pain and inflammation in osteoarthritis and rheumatoid arthritis
- Acute gout attacks
- Primary dysmenorrhoea
- Other conditions requiring rapid parenteral NSAID therapy

Dosage & Administration:

Adults:

- **Intramuscular:** 75 mg once daily, injected deeply into the gluteal muscle. In severe cases, 150 mg/day may be given as two separate injections 12 hours apart.
- **Intravenous infusion:** 75 mg diluted and infused over at least 30 minutes. Use the lowest effective dose for the shortest duration possible. Not recommended in children.

Contraindications:

- Hypersensitivity to diclofenac or other NSAIDs
- Active peptic ulcer or gastrointestinal bleeding
- History of asthma, urticaria, or acute rhinitis precipitated by NSAIDs
- Severe hepatic, renal, or cardiac failure

Warnings & Precautions:

- Use with caution in elderly patients

- Monitor for signs of gastrointestinal bleeding or ulceration
- Avoid in late pregnancy (third trimester)
- Use cautiously in patients with hypertension, heart disease, or impaired renal/hepatic function

Drug Interactions:

- Increased risk of bleeding with anticoagulants
- May increase serum lithium or digoxin levels
- Reduced efficacy of diuretics and antihypertensive drugs
- Concomitant NSAID use increases risk of adverse effects

Adverse Effects:

Common: Nausea, vomiting, abdominal pain, dyspepsia, dizziness, headache, injection site pain.
Serious (rare): Gastrointestinal bleeding, peptic ulcer, hepatotoxicity, severe skin reactions, anaphylaxis.

Overdose:

Symptoms: Drowsiness, vomiting, gastrointestinal bleeding, hypertension, renal failure, convulsions.

Treatment: Supportive and symptomatic. No specific antidote.

Storage:

Store below 25 °C in a dry place, protected from light. Do not freeze. Keep out of reach of children.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,
Panchkula-134113

TM: Trademark Applied for