

# Etastyle 250 Tablet

## Ethamsylate (250 mg) Tablet

---

### Generic Name:

Ethamsylate

### Strength:

250 mg

### Formulation:

Oral tablet

---

### Uses:

Ethamsylate is a **hemostatic agent** that helps control bleeding by improving blood clotting. It is primarily used for the following conditions:

- Menorrhagia (Heavy Menstrual Bleeding):**
    - Reduces excessive menstrual bleeding in women with menorrhagia.
  - Post-Surgical Bleeding:**
    - Minimizes bleeding after surgical procedures, especially in gynecological, dental, or other minor surgeries.
  - Capillary Bleeding:**
    - Controls bleeding caused by fragile blood vessels (capillaries), which may occur in conditions like **epistaxis** (nosebleeds).
  - Other Bleeding Disorders:**
    - Can be used for managing bleeding due to injuries or conditions affecting blood clotting.
- 

### Mechanism of Action:

Ethamsylate works by:

- Strengthening the Capillary Walls:** Increases the resistance and stability of small blood vessels to prevent bleeding.
  - Promoting Platelet Aggregation:** Enhances the ability of platelets to stick together and form clots at the site of bleeding.
  - Reducing Bleeding Time:** Shortens the time it takes for blood to clot and stop excessive bleeding.
- 

### Dosage:

- Typical Dose:**
    - 250 mg** tablet, taken 2-4 times daily, depending on the severity of the condition.
  - Take the tablet with a glass of water, with or without food.
  - Duration of Use:** Usually prescribed for short-term use, especially during active bleeding.
  - Follow your healthcare provider's instructions for exact dosage and duration.
- 

### Precautions:

- Pregnancy and Breastfeeding:**
  - Use only if prescribed by a healthcare professional after a careful assessment of risks and benefits.
- Kidney or Liver Disease:**
  - Caution is advised for individuals with kidney or liver disorders.
- Blood Clotting Disorders:**
  - Ethamsylate should be used with caution in individuals with bleeding or clotting disorders unless otherwise directed by a doctor.
- Alcohol Use:**

- Alcohol may affect the action of Ethamsylate, so it should be used with caution.
- 

## Side Effects:

- **Common:**
    - Nausea
    - Abdominal discomfort
    - Headache
  - **Less Common:**
    - Skin rashes
    - Dizziness
    - Diarrhea
  - **Serious (Seek Medical Attention):**
    - Severe allergic reactions (rash, itching, swelling of the face, lips, or throat)
    - Chest pain or difficulty breathing
    - Unexplained bruising or unusual bleeding
- 

## Drug Interactions:

Ethamsylate may interact with:

- **Anticoagulants** (e.g., warfarin)
- **Antiplatelet medications** (e.g., aspirin, clopidogrel)
- **Other bleeding medications**

Make sure to inform your doctor about all medications and supplements you are currently taking to avoid interactions.

---

## Storage:

- Store at **20°C to 25°C** (68°F to 77°F).
  - Keep the tablets in a **dry, cool place** away from light.
  - Keep out of reach of children.
- 

## Note:

Ethamsylate is intended for **short-term use** to manage and control bleeding. It should be used under the supervision of a healthcare provider, especially if you have underlying health conditions that affect bleeding or clotting. Always follow your healthcare provider's instructions carefully for the best results.

Manufactured in India for:

**Cafoli**<sup>TM</sup>  
L I F E C A R E

Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,  
Panchkula-134113

TM: Trademark Applied for