

Information For the Use
Ethamsylate (500mg)

Etastyle 500 Tablet

Ethamsylate (500 mg) Tablet

Generic Name:

Ethamsylate

Strength:

500 mg

Formulation:

Oral tablet

Uses:

Ethamsylate is a **hemostatic agent** used to control and prevent excessive bleeding in various conditions, including:

- Menorrhagia (Heavy Menstrual Bleeding):**
 - Reduces excessive blood loss during menstruation.
 - Post-Surgical Bleeding:**
 - Helps minimize bleeding after surgical procedures, especially in gynecological and dental surgeries.
 - Capillary Bleeding:**
 - Controls bleeding caused by fragile capillaries (small blood vessels).
 - Nosebleeds (Epistaxis):**
 - Effectively reduces bleeding from nasal passages.
 - Bleeding Due to Injury or Trauma:**
 - Aids in controlling blood loss caused by injuries.
-

Mechanism of Action:

Ethamsylate works by:

- Strengthening Capillaries:** Increases the resistance of blood vessel walls, reducing their permeability and tendency to bleed.
 - Promoting Platelet Adhesion:** Enhances the ability of platelets to adhere to damaged blood vessel walls, aiding in clot formation and reducing bleeding time.
 - Reducing Bleeding Time:** Helps the body form stable blood clots faster to stop bleeding effectively.
-

Dosage:

- Typical Dose:** 500 mg, 2 to 4 times a day, depending on the severity of the bleeding.
 - Route:** Oral, taken with a glass of water.
 - Follow your healthcare provider's recommendations for optimal results.
-

Precautions:

- Pregnancy and Breastfeeding:**
 - Use only if prescribed by a healthcare professional.
 - Blood Disorders:**
 - Inform your doctor if you have a history of clotting disorders.
 - Kidney or Liver Disease:**
 - Use with caution and under medical supervision.
 - Not for Use in:**
 - Severe bleeding disorders that require immediate surgical intervention.
 - Conditions where clot formation might be dangerous (e.g., recent heart attack, stroke).
-

Side Effects:

- Common:**
 - Nausea
 - Headache
 - Stomach pain

- Diarrhea
 - **Less Common:**
 - Skin rash or itching
 - Dizziness
 - **Serious (Seek Medical Attention):**
 - Severe allergic reactions (swelling of face, lips, or throat)
 - Shortness of breath
 - Chest pain
-

Drug Interactions:

- May interact with:
 - **Anticoagulants** (e.g., warfarin)
 - **Antiplatelet drugs** (e.g., aspirin, clopidogrel)

Inform your healthcare provider about all medications and supplements you are taking.

Storage:

- Store at **20°C to 25°C** (68°F to 77°F).
 - Protect from **light** and **moisture**.
 - Keep out of reach of children.
-

Note:

Ethamsylate is used for **short-term management of bleeding** and should be taken under medical supervision. It is not a replacement for surgical intervention when necessary. Follow your healthcare provider's instructions carefully for safe and effective use.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,
Panchkula-134113

TM: Trademark Applied for