# Salmeterol (25mcg) + Fluticasone Propionate (250mcg)

# Flutialfa S 250 Inhaler

Salmeterol (25 mcg) + Fluticasone Propionate (250 mcg) Inhaler

# **Description:**

This combination inhaler contains Salmeterol, a long-acting beta-agonist (LABA), and Fluticasone Propionate, a corticosteroid. It is used for the maintenance treatment of asthma and chronic obstructive pulmonary disease (COPD), providing long-term bronchodilation and anti-inflammatory action.

# **Composition:**

#### Each metered dose contains:

- Salmeterol Xinafoate: Equivalent to 25 mcg of Salmeterol
- Fluticasone Propionate: 250 mcg

#### Excipients: q.s.

#### **Indications:**

- Maintenance treatment of persistent asthma in adults and adolescents (12 years and older).
- Management of symptoms and prevention of exacerbations in COPD, including chronic bronchitis and emphysema.

# **Dosage and Administration:**

- Asthma and COPD:
  - 1 inhalation twice daily (morning and evening) or as directed by the physician.
- Rinse the mouth with water after each use to reduce the risk of oral thrush.

#### Note:

This inhaler is not intended for acute relief of bronchospasm. Use a short-acting bronchodilator (rescue inhaler) for sudden symptoms.

#### **Contraindications:**

- Hypersensitivity to Salmeterol, Fluticasone Propionate, or any component of the inhaler.
- Not for use in the treatment of acute bronchospasm or status asthmaticus.

# **Warnings and Precautions:**

- Asthma-related death: LABAs may increase the risk of asthma-related deaths if not used with an appropriate corticosteroid. Always use as a combination therapy.
- Regular monitoring is necessary for systemic corticosteroid effects, such as adrenal suppression or reduced bone mineral density, with long-term use.
- Use with caution in patients with cardiovascular disease, diabetes, thyroid disorders, or a history of seizures.
- Avoid abrupt discontinuation; tapering may be required under medical supervision.

#### **Side Effects:**

- Common: Headache, hoarseness, throat irritation, or tremors.
- Less common: Oral thrush, palpitations, or muscle cramps.

• Rare: Hypersensitivity reactions, paradoxical bronchospasm, or pneumonia (in COPD patients).

**Storage:** 

- Store at a temperature below 30°C (86°F).
- Keep the inhaler away from direct sunlight and heat.
- · Keep out of reach of children.

Note:

This inhaler is a maintenance medication and should be used regularly as prescribed. It is not a substitute for a rescue inhaler in acute situations. Always consult your healthcare provider for appropriate usage.

Manufactured in India for:



(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for