# Salmeterol (50mcg) + Fluticasone Propionate (250mcg)

# Flutialfa SF 250 Capsules

Salmeterol (50 mcg) + Fluticasone Propionate (250 mcg) AeroCaps

### **Description:**

Salmeterol is a long-acting beta-agonist (LABA) that provides sustained bronchodilation, and Fluticasone Propionate is an inhaled corticosteroid that reduces inflammation in the airways. The combination of these two medications in AeroCaps is used for the maintenance treatment of asthma and chronic obstructive pulmonary disease (COPD), helping to manage symptoms and prevent exacerbations.

## **Composition:**

# **Each capsule contains:**

- Salmeterol Xinafoate: Equivalent to 50 mcg of Salmeterol
- Fluticasone Propionate: 250 mcg

#### **Indications:**

- Maintenance treatment of persistent asthma in adults and adolescents (≥12 years).
- Long-term management of COPD to reduce symptoms and the frequency of exacerbations.

#### **Dosage and Administration:**

- Recommended Dose:
  - Asthma/COPD: Inhale the contents of 1 AeroCap twice daily (morning and evening) using the provided inhalation device.
- Instructions for Use:

- 1. Place the capsule in the inhalation device as per the instructions provided.
- 2. Pierce the capsule and inhale deeply through the mouthpiece.
- 3. Ensure the entire dose is inhaled.
- 4. Rinse the mouth with water after use to minimize the risk of oral thrush.

#### **Contraindications:**

- Hypersensitivity to Salmeterol, Fluticasone, or any components of the formulation.
- Not for use in the treatment of acute bronchospasm or status asthmaticus.

# **Warnings and Precautions:**

- Asthma-related deaths: LABAs may increase the risk of asthma-related deaths if not combined with corticosteroids. Use only as part of combination therapy.
- Regular monitoring is advised to detect potential systemic corticosteroid effects, such as adrenal suppression or reduced bone density, in long-term use.
- Use cautiously in patients with cardiovascular disorders, diabetes, thyroid dysfunction, or seizure disorders.
- Do not abruptly discontinue therapy; tapering under medical supervision may be necessary.

#### **Side Effects:**

- Common: Headache, throat irritation, dry mouth, or tremors.
- Less common: Oral thrush, hoarseness, or muscle cramps.

• Rare: Paradoxical bronchospasm, palpitations, or hypersensitivity reactions (rash, swelling, or difficulty breathing).

# **Storage:**

- Store below 25°C (77°F) in a dry place.
- Protect from light and moisture.
- · Keep out of reach of children.

#### **Note:**

This combination is a maintenance therapy and should not be used as a rescue treatment during acute asthma or COPD exacerbations. Always follow your healthcare provider's instructions regarding dosage and usage. Use a short-acting bronchodilator for sudden symptoms.

Manufactured in India for:



(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for