

Fusitrip Cream

Fusidic Acid 2% Cream

Composition:

Each gram of cream contains:

- **Fusidic Acid I.P.** – 2% w/w

Pharmacological Class:

- **Topical Antibiotic**

Indications:

Fusidic Acid 2% Cream is used for the treatment of **skin infections** caused by **Staphylococcus aureus** and other susceptible bacteria, including:

- **Impetigo**
- **Infected Dermatitis (Eczema, Atopic Dermatitis, Contact Dermatitis)**
- **Furunculosis (Boils)**
- **Infected Cuts, Wounds, and Burns**

Mechanism of Action:

Fusidic Acid is a **bacteriostatic antibiotic** that inhibits bacterial protein synthesis by blocking **elongation factor G (EF-G)**, preventing bacterial growth and spread.

Dosage & Administration:

- **Apply a thin layer** to the affected area **2-3 times daily** or as directed by a physician.
- **Do not cover with an occlusive dressing** unless prescribed.
- Treatment duration typically lasts **7-14 days**, depending on the severity of the infection.
- **Avoid contact with eyes, nose, and mouth.**

Contraindications:

- **Hypersensitivity** to Fusidic Acid or any of its excipients.
- Not suitable for **fungal or viral infections** (e.g., herpes, athlete's foot).

Warnings & Precautions:

- **Prolonged use may lead to antibiotic resistance.**
- **Avoid application on deep wounds or large areas** without medical supervision.
- **Do not use for more than 14 days** unless advised by a doctor.

Side Effects:

Common:

- Mild skin irritation (redness, itching, burning)
- Dry skin

Rare:

- **Allergic reactions** (rash, swelling, difficulty breathing – seek immediate medical attention)
- **Contact dermatitis**

Drug Interactions:

- Minimal systemic absorption, hence **low risk of interactions.**
- **Avoid using with other topical antibiotics** unless prescribed.

Storage Instructions:

- Store at **room temperature (below 30°C).**
- Keep away from direct sunlight and moisture.
- **Keep out of reach of children.**

Conclusion:

Fusidic Acid 2% Cream is a **potent topical antibiotic** for bacterial skin infections. It should be used **under medical supervision**, and prolonged use should be avoided to **prevent resistance development**.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,
Panchkula-134113

TM: Trademark Applied for