

Information For the User  
Gabapentin (6% w/w) + Lidocaine (5% w/w)

# Gabaxuro L Gel

## Gabaxuro L Gel

### Composition:

Each 100 g contains:

- **Gabapentin (6% w/w)**
- **Lidocaine (5% w/w)**

### Description:

**Gabaxuro L Gel** is a topical gel formulated with **Gabapentin** and **Lidocaine** designed for effective pain management. This combination gel provides targeted relief from localized pain, including neuropathic pain, muscle soreness, and post-surgical discomfort. Gabapentin, an anticonvulsant, works by stabilizing nerve activity, while **Lidocaine**, a local anesthetic, provides immediate numbing effect to relieve pain. Together, they offer a dual-action formula that not only reduces nerve-related pain but also numbs the area to alleviate discomfort quickly.

- **Gabapentin** targets nerve pain, preventing abnormal signals sent to the brain that cause pain sensations.
- **Lidocaine** blocks nerve signals in the skin, numbing the affected area and providing fast pain relief.

**Gabaxuro L Gel** is easy to apply, fast-absorbing, and non-greasy, making it ideal for use on a wide range of painful conditions.

### Pharmacological Action:

- **Gabapentin:** Works by inhibiting certain nerve signals that cause pain, making it particularly effective for neuropathic pain relief.
- **Lidocaine:** Temporarily numbs the application area by blocking nerve signal transmission, reducing discomfort and pain.

### Indications:

**Gabaxuro L Gel** is recommended for:

- Neuropathic pain (pain caused by nerve damage or dysfunction).
- Musculoskeletal pain (muscle strains, sprains, and stiffness).
- Post-surgical pain and localized discomfort.
- **Localized pain relief** for conditions such as sciatica, diabetic neuropathy, and shingles.

### Usage Instructions:

- **Recommended Dose:** Apply a thin layer of **Gabaxuro L Gel** to the affected area 2-3 times daily or as directed by your healthcare provider.
- **Application:** Gently massage the gel into the affected area until absorbed. Avoid applying to broken or irritated skin.
- For best results, use consistently over the prescribed period.

### Contraindications:

- Do not use if you are allergic to **Gabapentin**, **Lidocaine**, or any of the excipients in the product.
- Avoid use on open wounds or broken skin.
- Do not apply near the eyes, mucous membranes, or genital areas.

### Warnings & Precautions:

- For **external use only**.
- If skin irritation, redness, or rash occurs, discontinue use and consult a doctor.
- **Avoid contact with eyes.** In case of contact, rinse thoroughly with water.
- Use with caution in individuals with a history of **allergies to local anesthetics** or **neurological conditions**.
- If pain persists or worsens, consult a healthcare professional.

### Adverse Effects:

**Gabaxuro L Gel** is generally well-tolerated, but in some cases, may cause:

- Mild skin irritation or redness.
- Allergic reactions (rare).
- Burning or tingling sensations at the application site.
- Dryness or rash in the treated area.

### Drug Interactions:

- Inform your doctor if you are using other **topical medications** or have any ongoing medical treatments.
- Avoid using with other **local anesthetics** without medical advice.

### Storage Conditions:

- Store below **30°C** in a cool, dry place.
- Keep tightly closed and away from direct sunlight and moisture.
- Keep out of reach of children.

### Conclusion:

**Gabaxuro L Gel** offers fast and effective relief for localized pain, providing both nerve pain management and immediate numbing action. With **Gabapentin** and **Lidocaine**, it provides a dual-action formula for

those suffering from neuropathic pain, muscle pain, and post-surgical discomfort. Safe for topical use, **Gabaxuro L Gel** helps reduce pain while promoting comfort and healing for a variety of conditions.

Manufactured in India for:



**Cafoli Lifecare Pvt. Ltd.**

**(An ISO 9001: 2015 Certified Co.)**

Plot no.: 367-FF, Industrial Area Phase-I,  
Panchkula-134113

TM: Trademark Applied for