

Information For the User  
Gabapentin (8% w/w) + Diclofenac (4.64% w/w) + Methyl Salicylate (5% w/w)

# Gabaxuro Plus Gel

## Gabaxuro Plus Gel

### Composition:

Each 100 g contains:

- **Gabapentin (8% w/w)**
- **Diclofenac (4.64% w/w)**
- **Methyl Salicylate (5% w/w)**

### Description:

**Gabaxuro Plus Gel** is a powerful topical gel formulated with a combination of **Gabapentin**, **Diclofenac**, and **Methyl Salicylate** for effective pain relief. This triple-action formula targets both nerve pain and inflammation, offering fast and long-lasting relief from localized pain and discomfort.

- **Gabapentin (8% w/w)**: Works by modulating nerve activity, providing relief from neuropathic pain such as that caused by diabetes, shingles, or nerve injuries.
- **Diclofenac (4.64% w/w)**: A nonsteroidal anti-inflammatory drug (NSAID) that reduces inflammation and pain, commonly used for muscle, joint, and soft tissue injuries.
- **Methyl Salicylate (5% w/w)**: Provides a soothing, warming effect that enhances blood flow and further alleviates muscle pain and stiffness.

**Gabaxuro Plus Gel** is ideal for conditions such as arthritis, muscle strains, sprains, and general musculoskeletal pain, as well as nerve-related pain, making it a comprehensive solution for pain management.

### Pharmacological Action:

- **Gabapentin**: Inhibits abnormal nerve signaling and reduces neuropathic pain.
- **Diclofenac**: Reduces inflammation and pain associated with musculoskeletal disorders by inhibiting cyclooxygenase (COX) enzymes.
- **Methyl Salicylate**: Acts as a counterirritant to provide a cooling sensation that helps soothe and relax the affected area.

### Indications:

**Gabaxuro Plus Gel** is used for:

- **Neuropathic pain** (e.g., diabetic neuropathy, post-herpetic neuralgia).
- **Musculoskeletal pain** (e.g., muscle strains, sprains, and joint pain).
- **Arthritis pain** (e.g., osteoarthritis and rheumatoid arthritis).
- **Soft tissue injuries** (e.g., sprains, strains, and contusions).

### Dosage & Administration:

- **Recommended Dose:** Apply a thin layer of **Gabaxuro Plus Gel** to the affected area 2-3 times daily or as directed by your healthcare provider.
- Gently massage the gel into the skin until it is fully absorbed.
- Do not apply to broken or irritated skin.
- Wash hands thoroughly after application, unless the hands are the area being treated.

### Contraindications:

- **Gabaxuro Plus Gel** should not be used if you are allergic to **Gabapentin, Diclofenac, Methyl Salicylate**, or any of the excipients.
- Avoid applying the gel to broken, irritated, or infected skin.
- Do not use if you have a history of **aspirin-sensitive asthma** or **allergic reactions to NSAIDs**.
- Not recommended for use near eyes, mucous membranes, or genital areas.

### Warnings & Precautions:

- For **external use only**.
- Avoid contact with **eyes** or mucous membranes. In case of contact, rinse immediately with water.
- Use with caution in individuals with **liver or kidney disorders**, or a history of gastrointestinal bleeding.
- Not suitable for children under **12 years of age**.
- If **irritation** or allergic reaction occurs, discontinue use and consult a doctor.
- Consult your healthcare provider if symptoms persist or worsen.

### Adverse Effects:

**Gabaxuro Plus Gel** is generally well-tolerated, but in rare cases, it may cause:

- **Mild skin irritation** (redness, rash, or itching at the application site).
- **Burning or tingling** sensation in the treated area.
- **Dryness** or **peeling** of the skin.
- **Allergic reactions** (e.g., rash, swelling, or difficulty breathing in rare instances).

### Drug Interactions:

- **Gabaxuro Plus Gel** is unlikely to cause systemic drug interactions, as it is applied topically.
- However, inform your healthcare provider if you are using other topical medications or have existing medical conditions.
- Avoid using with other **topical NSAIDs** without medical advice.

### Storage Conditions:

- Store at **room temperature**, below **25°C**.
- Protect from moisture and direct sunlight.
- Keep out of reach of children.
- Close the cap tightly after use.

## Conclusion:

**Gabaxuro Plus Gel** provides a highly effective solution for relieving **localized pain** and **inflammation**. Combining **Gabapentin** for neuropathic pain, **Diclofenac** for inflammation, and **Methyl Salicylate** for a soothing warming effect, it offers comprehensive relief for conditions like **arthritis**, **musculoskeletal injuries**, and **nerve pain**. It is easy to apply, absorbs quickly, and provides fast-acting, long-lasting relief from pain and discomfort. Always use under the guidance of a healthcare professional for best results.

Manufactured in India for:



**Cafoli Lifecare Pvt. Ltd.**

**(An ISO 9001: 2015 Certified Co.)**

Plot no.: 367-FF, Industrial Area Phase-I,  
Panchkula-134113

TM: Trademark Applied for