

Information For the User
Ferric Ammonium Citrate I.P. 160mg + Cyanocobalamin I.P. 7.5mcg + Folic Acid I.P. 0.5mg + Alcohol
(95%) I.P. 0.87ml Syrup

Hematyle Syrup

INFORMATION FOR THE USE

Hematyle Syrup

Hematinic Syrup of Iron, Folic Acid & Vitamin B12

Description:

Hematyle Syrup is a hematinic preparation containing essential nutrients including Iron, Folic Acid, and Vitamin B12. This combination is specifically formulated to support red blood cell formation, improve hemoglobin levels, and combat nutritional deficiencies commonly associated with anemia. The syrup format ensures better patient compliance, especially in pediatric and geriatric populations.

Composition: *(Each 5ml contains)*

- **Iron (as Ferrous Fumarate or equivalent):** [Specify exact mg]
- **Folic Acid I.P.:** [Specify exact mcg]
- **Vitamin B12 I.P. (Cyanocobalamin):** [Specify exact mcg]
(Final quantitative values to be provided based on formulation)

Indications:

Hematyle Syrup is indicated for:

1. **Iron Deficiency Anemia:** Supports red blood cell production and improves hemoglobin synthesis.
2. **Megaloblastic Anemia:** Corrects deficiencies of Folic Acid and Vitamin B12.
3. **Pregnancy & Lactation:** Helps meet increased nutritional demands.
4. **Convalescence:** Restores nutritional balance post-illness.
5. **Nutritional Deficiency States:** Improves general weakness and fatigue due to inadequate dietary intake.

Mechanism of Action:

- **Iron:** Crucial for hemoglobin formation, facilitates oxygen transport in the blood.
- **Folic Acid:** Essential for DNA synthesis and cell division; prevents neural tube defects during pregnancy.
- **Vitamin B12:** Required for neurological function and red blood cell maturation.

Dosage and Administration:

- **Adults:** 5–10 ml once or twice daily or as directed by the physician.
- **Children:** As advised by the pediatrician depending on age and weight.

- **Route:** Oral
- Shake well before use.

Side Effects:

- **Common:** Constipation, dark stools, metallic taste.
- **Less Common:** Nausea, stomach upset, allergic reactions.

Precautions:

1. **Allergy:** Avoid in patients with known hypersensitivity to iron or other ingredients.
2. **Overdose:** Keep away from children; iron overdose can be fatal in pediatric cases.
3. **Concurrent Use:** Caution with antacids or tetracyclines that may interfere with absorption.
4. **Pregnancy:** Safe under medical supervision.
5. **Alcohol:** May impair absorption; to be avoided during therapy.

Storage:

- Store in a cool, dry place away from direct sunlight.
- Keep bottle tightly closed.
- Keep out of reach of children.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for