

Levocetirizine (2.5mg) + Montelukast (4mg) Bilayered

Monclip LC Kid Tablet

**Levocetirizine (2.5 mg) + Montelukast (4 mg)
Bilayered Tablet**

Description:

This bilayered tablet combines Levocetirizine, a second-generation antihistamine, and Montelukast, a leukotriene receptor antagonist. It is used to manage symptoms of allergic rhinitis and asthma by blocking histamine and leukotrienes, reducing inflammation and allergic reactions.

Composition (Each Bilayered Tablet Contains):

- Levocetirizine Dihydrochloride: 2.5 mg
- Montelukast Sodium equivalent to Montelukast: 4 mg
- Excipients: q.s.

Indications:

- Allergic rhinitis (seasonal and perennial)
- Hay fever
- Allergic asthma
- Relief of symptoms such as:
 - Sneezing
 - Runny nose
 - Nasal congestion
 - Watery, itchy eyes
 - Breathing difficulties

Dosage and Administration:

Recommended Dosage:

- Children (2 to 6 years):
 - One tablet daily in the evening, as directed by the physician.

Route of Administration:

- Oral. Swallow the tablet with water. Do not crush or chew.

Contraindications:

- Hypersensitivity to Levocetirizine, Montelukast, or any excipients.
- Severe renal impairment (creatinine clearance <10 mL/min).

Warnings and Precautions:

1. Neuropsychiatric Events:
 - Montelukast has been associated with behavioral changes, agitation, anxiety, and rare suicidal ideation. Monitor patients for such symptoms.
2. Sedation:
 - Levocetirizine may cause mild drowsiness; caution is advised when engaging in activities requiring alertness.
3. Asthma Control:
 - Not suitable for acute asthma attacks; continue prescribed rescue medications.
4. Renal Impairment:
 - Dose adjustments may be required for patients with reduced kidney function.

Side Effects:

- Common Side Effects:
 - Headache
 - Drowsiness or fatigue
 - Dry mouth
 - Abdominal pain
 - Nausea
- Serious Side Effects:

- Allergic reactions (e.g., rash, swelling, difficulty breathing)
- Behavioral changes (associated with Montelukast)
- Severe drowsiness

Note:

This is a prescription-only medication. Use under the guidance of a healthcare provider and adhere to the prescribed dosage.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for

Drug Interactions:

- **Alcohol and CNS Depressants:**
 - May enhance sedative effects of **Levocetirizine. Avoid concurrent use.**
- **CYP450 Inducers:**
 - Drugs like rifampin may reduce Montelukast levels.
- **Other Antihistamines:**
 - Avoid simultaneous use to prevent additive sedation.

Use in Specific Populations:

- **Pregnancy:**
 - Use only if clearly indicated and prescribed by a healthcare provider.
- **Lactation:**
 - Caution is advised as it is unknown whether the components are excreted in breast milk.
- **Pediatrics:**
 - Safe for children aged 2 to 6 years under medical supervision.
- **Geriatrics:**
 - Monitor for renal function and potential side effects; no specific dose adjustments needed unless renal impairment is present.

Storage Instructions:

- Store at a temperature of 15°C to 30°C (59°F to 86°F).
 - Protect from light and moisture.
 - Keep out of reach of children.
-