

Moxolunic BF Eye Drop

**Bromfenac (0.09% w/v) + Moxifloxacin (0.5% w/v)
Eye Drops**

Drug Class:

- **Bromfenac: Non-Steroidal Anti-Inflammatory Drug (NSAID)**
- **Moxifloxacin: Fluoroquinolone Antibiotic**

Description:

This combination eye drop contains Bromfenac, an anti-inflammatory agent that reduces pain and swelling, and Moxifloxacin, an antibiotic that treats bacterial infections. Together, they are used to manage post-operative eye inflammation and prevent or treat bacterial infections in the eyes.

Indications:

- **Treatment of post-operative inflammation and pain following eye surgeries such as cataract surgery.**
- **Prevention and treatment of bacterial infections in the eye.**

Dosage and Administration:

- **Dosage:** Instill 1 drop in the affected eye(s) as directed by your doctor, typically 2–4 times daily or as prescribed.
- **Administration:**
 1. Wash your hands thoroughly before using the eye drops.
 2. Tilt your head back slightly and gently pull down the lower eyelid to create a small pocket.
 3. Instill one drop into the pocket, then close your eyes for 1–2 minutes.
 4. Avoid touching the dropper tip to any surface, including your eye, to prevent contamination.

Contraindications:

- **Hypersensitivity:** Do not use if allergic to Bromfenac, Moxifloxacin, or any other ingredients in the formulation.
- **Pregnancy and breastfeeding:** Should be used only if clearly needed and under medical supervision.

Warnings and Precautions:

- **Ocular sensitivity:** May cause temporary stinging or discomfort upon application.
- **Use with contact lenses:** Avoid wearing contact lenses while using the drops or if signs of infection are present.
- **Delayed healing:** Bromfenac may delay wound healing. Use cautiously in patients with corneal disorders.
- **Avoid long-term use:** Prolonged use may lead to increased intraocular pressure or corneal complications.

Side Effects:

Common side effects include:

- Temporary stinging, burning, or itching of the eyes
- Blurred vision immediately after application
- Dry or irritated eyes

Rare but serious side effects include:

- Severe eye pain or redness
- Vision changes
- Allergic reactions (e.g., swelling, rash, or difficulty breathing)

Drug Interactions:

- **Other topical eye medications:** If using multiple eye drops, wait at least 5–10 minutes between applications.
 - **NSAIDs and corticosteroids:** Concurrent use may increase the risk of corneal complications.
-

Storage:

- **Store in a cool, dry place, below 25°C.**
 - **Protect from light and moisture.**
 - **Keep out of reach of children.**
 - **Do not use after the expiry date or if the solution changes color.**
-

Conclusion:

Bromfenac + Moxifloxacin eye drops provide dual benefits by reducing inflammation and preventing bacterial infections. Always use as prescribed by your doctor, and report any unusual symptoms or side effects. Regular follow-up is recommended for monitoring response to treatment.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,
Panchkula-134113

TM: Trademark Applied for