

Nepasil Eye Drop

Nepasil Eye Drop (Nepafenac 0.1% w/v)

Introduction

Nepasil Eye Drop contains **Nepafenac 0.1% w/v**, a **nonsteroidal anti-inflammatory drug (NSAID)** used to reduce **pain, inflammation, and swelling** in the eyes, particularly after **cataract surgery**. It helps manage **post-operative eye discomfort** and is also effective in treating **ocular pain and inflammation caused by various conditions**.

Composition & Mechanism of Action

1. **Nepafenac (0.1% w/v)**
 - A **prodrug of Amfenac**, which gets converted into its active form in the eye.
 - **Inhibits cyclooxygenase (COX-1 & COX-2) enzymes**, reducing the production of **prostaglandins** that cause **pain and inflammation**.
 - Provides **long-lasting relief from post-surgical inflammation, swelling, and redness**.

Indications & Uses

Nepasil Eye Drop is used for:

- **Post-operative inflammation and pain** after **cataract surgery**.
- **Ocular pain, swelling, and redness** due to eye procedures.
- **Treatment of eye inflammation** associated with various eye disorders.

Dosage & Administration

- **Recommended Dosage:**
 - **Before Surgery:** 1 drop in the affected eye **three times a day** starting **one day before surgery**.
 - **Post-Surgery:** Continue using for **2-4 weeks** as directed by a doctor.
- **How to Use:**
 - Wash hands before application.
 - Tilt the head back and pull down the lower eyelid to create a small pocket.
 - Instill **one drop into the affected eye** and close the eye for a few seconds.
 - Avoid touching the dropper tip to prevent contamination.

Precautions & Warnings

- **Not recommended** for patients allergic to **NSAIDs or Nepafenac**.
- **Avoid use in pregnant or breastfeeding women** unless prescribed by a doctor.
- **Do not wear contact lenses** while using this medication.
- Use with **caution in diabetic patients** or those with **corneal conditions**.

- May cause **temporary blurred vision** after application—avoid driving or operating machinery immediately after use.

Side Effects

- **Common Side Effects:**
 - Mild **eye irritation or burning**
 - Temporary **blurry vision**
 - **Dry eyes or watery eyes**
- **Serious Side Effects (Rare):**
 - **Eye pain or swelling**
 - **Increased sensitivity to light**
 - **Signs of infection (eye redness, discharge, or worsening inflammation)**

Storage & Handling

- Store at **15°C – 25°C** in a **cool, dry place**, away from direct sunlight.
- Keep the bottle **tightly closed** and do not freeze.
- Discard the bottle **one month after opening**.

Conclusion

Nepasil Eye Drop (Nepafenac 0.1% w/v) is a **trusted ophthalmic NSAID** used for **managing post-surgical pain and inflammation** in the eyes. With its **potent anti-inflammatory action**, it provides **quick relief from discomfort and supports faster healing** after cataract surgery and other eye procedures.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for