Ofloglee 200 Tablet

Ofloxacin 200 mg Tablet

Description:

Ofloxacin 200 mg Tablet is a broadspectrum antibiotic used to treat various bacterial infections. It belongs to the fluoroquinolone class of antibiotics and works by inhibiting bacterial DNA replication, effectively killing the bacteria responsible for the infection.

Composition:

• Ofloxacin: 200 mg

Indications:

This medication is indicated for the treatment of bacterial infections such as:

- 1. Respiratory Tract Infections: Bronchitis, pneumonia, and sinusitis.
- 2. Urinary Tract Infections (UTIs): Including cystitis and pyelonephritis.
- 3. Gastrointestinal Infections: Traveler's diarrhea, typhoid fever, and other intestinal infections.
- 4. Skin and Soft Tissue Infections: Boils, abscesses, and infected wounds.
- 5. Sexually Transmitted Infections: Gonorrhea and pelvic inflammatory disease.

Mechanism of Action:

Ofloxacin inhibits bacterial DNA gyrase and topoisomerase IV, enzymes critical for DNA replication and repair in bacteria. This prevents the bacteria from multiplying and eventually leads to their elimination.

Dosage and Administration:

- Usual Adult Dose: 200–400 mg per day, depending on the severity and type of infection.
- Route: Oral, taken with or without food.
- Always follow the prescribed dosage by your doctor.

Side Effects:

- Common: Nausea, vomiting, diarrhea, dizziness, or headache.
- Rare but Serious: Tendon rupture, QT prolongation, allergic reactions (rash, swelling, difficulty breathing).
- Stop use and seek medical attention if severe side effects occur.

Precautions:

- 1. Avoid excessive exposure to sunlight or UV light; Ofloxacin can cause photosensitivity.
- 2. Use with caution in individuals with kidney disorders, epilepsy, or heart rhythm issues.

- 3. Avoid concurrent use with antacids, calcium, iron, or magnesium supplements, as they may reduce absorption.
- 4. Not recommended for children, adolescents, or pregnant/breastfeeding women unless prescribed by a doctor.
- 5. Avoid alcohol as it may enhance side effects like dizziness.

Storage:

- Store at room temperature (15-30°C) in a dry place away from light and moisture.
- · Keep out of reach of children.

Note: This medication should only be used under medical supervision. Completing the full prescribed course is essential, even if symptoms improve, to prevent antibiotic resistance.

Manufactured in India for:



(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for