

Panpital 20 Tablet

Pantoprazole (20mg) Tablet

antibiotics to eliminate *H. pylori* infection.

Description

Pantoprazole (20mg) Tablet is a proton pump inhibitor (PPI) that effectively reduces stomach acid production. It is used to manage conditions such as acid reflux, peptic ulcers, and other disorders caused by excess stomach acid. Pantoprazole works by inhibiting the proton pumps in the stomach lining, which are responsible for acid secretion.

Composition

- **Active Ingredient:**
 - **Pantoprazole Sodium Sesquihydrate (equivalent to Pantoprazole): 20mg**

Indications

This medication is indicated for the treatment of:

- **Gastroesophageal Reflux Disease (GERD):** Provides relief from heartburn and acid regurgitation.
- **Erosive Esophagitis:** Promotes healing of damage to the esophageal lining caused by stomach acid.
- **Peptic Ulcers:** Aids in healing stomach and duodenal ulcers and prevents recurrence.
- **Zollinger-Ellison Syndrome:** Manages excessive acid production caused by gastrin-secreting tumors.
- **Helicobacter pylori Eradication Therapy:** Used in combination with

Dosage and Administration

- **Recommended Dosage:**
 - **Adults:** One tablet (20mg) once daily, usually before breakfast.
 - **Duration:** Treatment duration depends on the condition and should be as prescribed by a healthcare provider.
- **Administration:**
 - Swallow the tablet whole with water. Do not crush, chew, or break the tablet.

Note: Always follow your doctor's instructions for dosage and duration.

Contraindications

- **Hypersensitivity to Pantoprazole or any other proton pump inhibitors.**
- **Severe liver impairment:** Use with caution and under medical supervision.
- **Not recommended for use in children under 12 years unless prescribed by a doctor.**

Warnings and Precautions

- **Long-Term Use:** Prolonged use of PPIs may increase the risk of bone fractures, vitamin B12 deficiency, and low magnesium levels.
- **Rebound Acid Hypersecretion:** Discontinuing abruptly may lead to

increased stomach acid. Tapering off is recommended.

- **Clostridium difficile Infection:** Long-term use may increase the risk of gut infections.
- **Liver Function:** Monitor liver enzymes in patients with a history of liver disease.
- **Pregnancy and Lactation:** Use only if clearly needed and prescribed by a healthcare provider.

- **Digoxin:** Increased risk of digoxin toxicity; monitor serum levels.
- **Methotrexate:** May elevate methotrexate levels; use cautiously in high doses.

Inform your doctor about all medications you are currently taking to avoid potential interactions.

Adverse Effects

Common Side Effects:

- Headache
- Nausea or vomiting
- Diarrhea or constipation
- Abdominal pain

Rare but Serious Side Effects:

- Severe allergic reactions (rash, swelling, difficulty breathing)
- Low magnesium levels (symptoms include muscle spasms, irregular heartbeat, or seizures)
- Kidney issues (e.g., changes in urine output, swelling)

If severe side effects occur, discontinue use and consult a doctor immediately.

Storage

- Store at room temperature (15°C to 30°C), away from moisture and direct sunlight.
- Keep out of reach of children.

Key Points

- **Effective Acid Control:** Quickly reduces stomach acid and promotes healing of acid-related conditions.
- **Once-Daily Dosage:** Convenient dosing regimen for GERD and related conditions.

Note: This medication should only be used as prescribed by a healthcare professional. Report any unusual symptoms or side effects to your doctor promptly.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,
Panchkula-134113

TM: Trademark Applied for

Drug Interactions

- **Antifungal Agents:** May reduce the effectiveness of ketoconazole and itraconazole.
- **Clopidogrel:** May reduce the efficacy of this antiplatelet drug.
- **Warfarin:** Increased risk of bleeding; monitor INR levels.