

Panpital DSR Capsule

Domperidone (30 mg) + Pantoprazole (40 mg) Capsule

Description:

Domperidone 30 mg and Pantoprazole 40 mg capsule is a combination medication used to treat various gastrointestinal conditions related to acid reflux, delayed gastric emptying, and gastric ulcers. Domperidone is a prokinetic agent that enhances gastric motility, while Pantoprazole is a proton pump inhibitor (PPI) that reduces the production of stomach acid. Together, they help in managing symptoms like nausea, bloating, acid reflux, and indigestion.

Composition:

- Domperidone: 30 mg
 - Pantoprazole: 40 mg
-

Indications:

This combination capsule is indicated for the treatment of:

1. **Gastroesophageal Reflux Disease (GERD):** To control acid reflux, heartburn, and regurgitation.
2. **Gastric Ulcers and Duodenal Ulcers:** To treat and prevent the recurrence of ulcers by reducing stomach acid secretion.

3. **Dyspepsia (Indigestion):** Helps in alleviating discomfort caused by acid in the stomach.
 4. **Nausea and Vomiting:** Reduces symptoms of nausea and vomiting, especially in conditions related to acid reflux.
 5. **Gastroparesis:** Improves stomach motility and alleviates symptoms related to delayed gastric emptying.
-

Mechanism of Action:

- **Domperidone:** Acts as a dopamine antagonist, increasing the motility of the stomach and intestines, facilitating faster gastric emptying, and preventing nausea and vomiting.
 - **Pantoprazole:** Inhibits the proton pump (H⁺/K⁺ ATPase) in the stomach lining, reducing the production of gastric acid and providing relief from acid-related conditions.
-

Dosage and Administration:

- **Usual Adult Dose:**
 - One capsule (30 mg Domperidone + 40 mg Pantoprazole) taken once daily, preferably 30 minutes before a meal, or as directed by a healthcare provider.
- **Pediatric Use:** Consult a healthcare provider before use in children.
- **Administration:**

- Swallow the capsule whole with a glass of water. Do not crush or chew the capsule.

- 4. **Pregnancy and Breastfeeding:** Consult a healthcare provider before using this medication during pregnancy or breastfeeding.
- 5. **Alcohol:** Avoid excessive alcohol consumption, as it can exacerbate gastrointestinal symptoms and interact with the medications.

Side Effects:

- **Common:**
 - Diarrhea or constipation, headache, dizziness, abdominal discomfort, dry mouth, mild nausea, or changes in taste sensation.
- **Serious but Rare:**
 - **Cardiac Effects:** Irregular heartbeats or QT prolongation (especially in patients with existing heart conditions).
 - **Endocrine Effects:** Gynecomastia (enlarged breast tissue in men) or lactation.
 - **Allergic Reactions:** Rash, itching, swelling, or difficulty breathing.
 - **Severe Gastrointestinal Reactions:** Persistent stomach pain, black stools, or signs of gastrointestinal bleeding.

Storage:

- Store at room temperature (15-30°C), away from direct sunlight, heat, and moisture.
- Keep the capsules tightly closed and out of reach of children.

Note: Domperidone + Pantoprazole capsules should be used according to your healthcare provider's instructions. Avoid exceeding the recommended dose. If symptoms persist or worsen, consult your doctor for further advice. If you experience any side effects, particularly severe reactions like chest pain, palpitations, or allergic symptoms, seek medical attention immediately.

Precautions:

1. **Liver Disease:** Use with caution in patients with liver impairment, as Pantoprazole is metabolized in the liver.
2. **Heart Conditions:** Patients with a history of cardiac arrhythmias or prolonged QT intervals should be closely monitored.
3. **Kidney Function:** Renal impairment may alter the metabolism of Domperidone.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for