# Information For the User Telmisartan 80mg + Hydrochlorothiazide 12.5mg Tablet

# **Telbolife H 80 Tablet**

**Brand Name: Telbolife H 80 Tablet** 

**Dosage Form**: Tablet

Therapeutic Category: Antihypertensive Combination

## **Composition:**

Each tablet contains:

• **Telmisartan I.P.** – 80 mg

• **Hydrochlorothiazide I.P.** – 12.5 mg

## **Description**:

**Telbolife H 80 Tablet** is a combination of **Telmisartan**, an **angiotensin II receptor blocker** (**ARB**), and **Hydrochlorothiazide**, a **thiazide diuretic**, used for the treatment of **hypertension**.

- **Telmisartan** works by blocking the action of **angiotensin II**, resulting in relaxation of blood vessels and a reduction in blood pressure.
- **Hydrochlorothiazide** helps the body eliminate excess sodium and water, which lowers blood pressure by reducing fluid volume.

#### **Indications:**

- **Essential Hypertension**: For the treatment of **high blood pressure** (hypertension).
- **Prevention of Stroke and Cardiovascular Events**: Reduces the risk of **stroke** and other cardiovascular complications in hypertensive patients.

#### **Dosage and Administration:**

- **Recommended Dose**: 1 tablet once daily, preferably in the morning.
- **Administration**: The tablet can be taken with or without food. It is advised to take it consistently at the same time each day for maximum benefit.
- **Dose Adjustment**: The dosage may be adjusted based on individual response and clinical assessment.

#### **Contraindications:**

- **Hypersensitivity** to any component of the formulation.
- Severe renal impairment or anuria.
- Hypersensitivity to sulfonamide derivatives (due to Hydrochlorothiazide).
- Pregnancy and breastfeeding.

#### **Precautions:**

• **Renal Function**: Regular monitoring of kidney function is recommended in patients with pre-existing renal impairment.

- **Electrolyte Imbalance**: Monitor **potassium** and **sodium** levels due to the diuretic effects of **Hydrochlorothiazide**.
- **Volume Depletion**: Caution is advised in patients with volume depletion or those on a low-salt diet.
- **Liver Disease**: Use with caution in patients with liver dysfunction, and monitor liver function as needed.

#### **Adverse Effects**:

- Common Side Effects: Dizziness, headache, fatigue, and gastrointestinal discomfort.
- **Serious Adverse Effects**: Renal dysfunction, **electrolyte imbalances** (especially **hypokalemia**), hypotension, and dehydration.
- Other Effects: Rash, muscle cramps, and elevated blood sugar levels.

#### **Storage**:

- Store below 25°C in a cool, dry place.
- Protect from moisture and direct sunlight.
- Keep out of the reach of children.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for