

Uriviga F MR Tablet

Tamsulosin 0.4 mg + Finasteride 5 mg Tablet

Dual-Action Therapy for Benign Prostatic Hyperplasia (BPH)

This combination tablet of Tamsulosin and Finasteride is designed to provide effective relief from symptoms of Benign Prostatic Hyperplasia (BPH). By combining a selective alpha-blocker and a 5-alpha reductase inhibitor, it addresses both the immediate and long-term management of BPH, helping improve urinary flow, reduce prostate size, and enhance the quality of life for men suffering from this condition.

Composition & Mechanism of Action

- Tamsulosin (0.4 mg)**
 - Class:** Alpha-1 adrenergic receptor blocker.
 - Action:** Relaxes the smooth muscles of the bladder neck and prostate, improving urine flow and reducing symptoms of urinary obstruction like frequent urination and difficulty in starting urination.
- Finasteride (5 mg)**
 - Class:** 5-alpha reductase inhibitor.
 - Action:** Reduces the size of the prostate gland by inhibiting the conversion of testosterone to dihydrotestosterone (DHT), which is a key factor in prostate enlargement.

Key Benefits

- Improves Urinary Flow:** Relieves symptoms such as weak stream, frequent urination, and urgency.
- Reduces Prostate Size:** Helps shrink the prostate over time, addressing the root cause of BPH.

- Dual Action:** Combines immediate symptomatic relief (Tamsulosin) with long-term treatment benefits (Finasteride).
- Enhances Quality of Life:** Reduces discomfort associated with BPH and improves bladder control.

Indications

- Benign Prostatic Hyperplasia (BPH) in adult men.
- Management of lower urinary tract symptoms associated with BPH.

Dosage and Administration

- Recommended Dose:** One tablet daily or as directed by a healthcare professional.
- Swallow the tablet whole with water, preferably at the same time each day.
- Can be taken with or without food.

Precautions

- Not for Use by Women or Children:** This medication is specifically for adult men and should not be handled by pregnant women, as Finasteride can harm a developing male fetus.
- Monitor Blood Pressure:** Tamsulosin may cause a drop in blood pressure, especially upon standing.
- Prostate Cancer Screening:** Rule out prostate cancer before starting treatment, as symptoms may overlap.
- Allergic Reactions:** Discontinue use and seek medical advice if symptoms like rash, itching, or swelling occur.

Potential Side Effects

- **Common:** Dizziness, headache, low blood pressure, nasal congestion, and ejaculation disorders.
 - **Less Common:** Breast tenderness or enlargement, reduced libido, and impotence (associated with Finasteride).
 - **Rare:** Severe allergic reactions or angioedema.
-

Additional Information

- **Storage:** Keep in a cool, dry place away from direct sunlight.
- **Packaging:** Available in blister packs of 10 or 30 tablets.
- **Shelf Life:** Best before 24 months from the date of manufacture.

Take control of BPH symptoms with Tamsulosin + Finasteride Tablets. A trusted dual-action solution for improved urinary health and prostate management. Consult your healthcare provider for tailored advice and care.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,
Panchkula-134113

TM: Trademark Applied for